

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12Q 1/68, G01N 33/53, 33/574		A1	(11) International Publication Number: WO 99/60161
		(43) International Publication Date: 25 November 1999 (25.11.99)	
(21) International Application Number: PCT/US99/10498		(74) Agents: LICATA, Jane, Massey et al.; Law Offices of Jane Massey Licata, 66 E. Main Street, Marlton, NJ 08053 (US).	
(22) International Filing Date: 12 May 1999 (12.05.99)		(81) Designated States: CA, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(30) Priority Data: 60/086,266 21 May 1998 (21.05.98) US		Published <i>With international search report.</i>	
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 60/086,266 (CIP) Filed on 21 May 1998 (21.05.98)			
(71) Applicant (for all designated States except US): DIADEXUS LLC [US/US]; 3303 Octavius Drive, Santa Clara, CA 95054 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): MACINA, Roberto, A. [AR/US]; 4118 Crescendo Avenue, San Jose, CA 95136 (US). YANG, Fei [CN/US]; Apartment 204, 18375 Caminito Cantilena, San Diego, CA 92128 (US). SUN, Yongming [CN/US]; Apartment 260, 869 S. Winchester Boulevard, San Jose, CA 92128 (US).			
(54) Title: A NOVEL METHOD OF DIAGNOSING, MONITORING, AND STAGING COLON CANCER			
(57) Abstract The present invention provides a new method for detecting, diagnosing, monitoring, staging, and prognosticating colon cancer.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**A NOVEL METHOD OF DIAGNOSING,
MONITORING, AND STAGING COLON CANCER**

FIELD OF THE INVENTION

This invention relates, in part, to newly developed assays for detecting, diagnosing, monitoring, staging, and prognosticating cancers, particularly colon cancer.

BACKGROUND OF THE INVENTION

Colon cancer is the second most frequently diagnosed malignancy in the United States. Cancer of the gastrointestinal tract, especially colon cancer, is a highly treatable and often a curable disease when localized to the bowel. However, currently colon cancer is the second most common cause of cancer death. Surgery is the primary treatment and results in cure in approximately 50% of patients. Recurrence following surgery is a major problem and often is the ultimate cause of death. The prognosis of colon cancer is clearly related to the degree of penetration of the tumor through the bowel wall and the presence or absence of nodal involvement. These two characteristics form the basis for all staging systems developed for this disease. Bowel obstruction and bowel perforation are indicators of poor prognosis. Elevated pretreatment serum levels of carcinoembryonic antigen (CEA) and carbohydrate antigen 19-9 (CA 19-9) also have negative prognostic significance.

Because of the frequency of the disease (approximately 160,000 new cases of colon cancer per year), the identification of high-risk groups, the demonstrated slow growth of primary lesions, the better survival of early-stage lesions, and the relative simplicity and accuracy of screening tests, screening for colon cancer should be a part of routine

care for all adults starting at age 50, especially those with first-degree relatives with colorectal cancer.

Procedures used for detecting, diagnosing, monitoring, staging, and prognosticating colon cancer are of critical importance to the outcome of the patient. For example, patients diagnosed with early colon cancer generally have a much greater five-year survival rate as compared to the survival rate for patients diagnosed with distant metastasized colon cancer. Treatment decisions are usually made in reference to the older Dukes or the Modified Astler-Coller (MAC) classification schema for staging. However, new diagnostic methods which are more sensitive and specific for detecting early colon cancer are clearly needed.

Further, colon cancer patients must be closely monitored following initial therapy and during adjuvant therapy to determine response to therapy and to detect persistent or recurrent disease or metastasis. Thus, there is clearly a need for a colon cancer marker which is more sensitive and specific in detecting colon cancer recurrence.

Another important step in managing colon cancer is to determine the stage of the patient's disease. Stage determination has potential prognostic value and provides criteria for designing optimal therapy. Currently, pathological staging of colon cancer is preferable over clinical staging as pathological staging provides a more accurate prognosis. However, clinical staging would be preferred were the method of clinical staging at least as accurate as pathological staging because it does not depend on an invasive procedure to obtain tissue for pathological evaluation. Staging of colon cancer would be improved by detecting new markers in cells, tissues, or bodily fluids which could differentiate between different stages of invasion.

In the present invention, methods are provided for detecting, diagnosing, monitoring, staging, and

- 3 -

prognosticating colon cancers, particularly colon, stomach, and small intestine cancer, via nine (9) Colon Specific Genes (CSGs). The nine CSGs refer, among other things, to native proteins expressed by the genes comprising the polynucleotide sequences of any of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8 or 9. In the alternative, what is meant by the nine CSGs as used herein, means the native mRNAs encoded by the genes comprising any of the polynucleotide sequences of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8 or 9 or levels of the genes comprising any of the polynucleotide sequences of SEQ ID NO:1, 2, 3, 4, 5, 6, 7, 8 or 9.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

SUMMARY OF THE INVENTION

Toward these ends, and others, it is an object of the present invention to provide a method for diagnosing the presence of colon cancer in a patient which comprises measuring levels of CSG in a sample of cells, tissue or bodily fluid from the patient and comparing the measured levels of CSG with levels of CSG in preferably the same cells, tissue, or bodily fluid type of a control, wherein an increase in the measured CSG levels in the patient versus levels of CSG in the control is associated with colon cancer.

Another object of the present invention is to provide a method of diagnosing metastatic colon cancer in a patient which comprises measuring CSG levels in a sample of cells,

- 4 -

tissue, or bodily fluid from the patient and comparing the measured CSG levels with levels of CSG in preferably the same cells, tissue, or bodily fluid type of a control, wherein an increase in measured CSG levels in the patient versus levels of CSG in the control is associated with a cancer which has metastasized.

Another object of the present invention is to provide a method of staging colon cancer in a patient which comprises identifying a patient having colon cancer, measuring levels of CSG in a sample of cells, tissues, or bodily fluid obtained from the patient, and comparing the measured CSG levels with levels of CSG in preferably the same cells, tissue or bodily fluid type of a control. An increase in measured CSG levels in the patient versus CSG levels in the control can be associated with a cancer which is progressing while a decrease or equivalent level of CSG measured in the patient versus the control can be associated with a cancer which is regressing or in remission.

Another object of the present invention is to provide a method of monitoring colon cancer in a patient for the onset of metastasis. The method comprises identifying a patient having colon cancer that is not known to have metastasized, periodically measuring levels of CSG in a sample of cells, tissues, or bodily fluid obtained from the patient, and comparing the measured CSG levels with levels of CSG in preferably the same cells, tissue, or bodily fluid type of a control, wherein an increase in measured CSG levels versus control CSG levels is associated with a cancer which has metastasized.

Yet another object of the present invention is to provide a method of monitoring the change in stage of colon cancer in a patient which comprises identifying a patient having colon cancer, periodically measuring levels of CSG in a sample of cells, tissue, or bodily fluid obtained from the patient, and comparing the measured CSG levels with levels of

- 5 -

CSG in preferably the same cells, tissues, or bodily fluid type of a control wherein an increase in measured CSG levels versus the control CSG levels is associated with a cancer which is progressing and a decrease in the measured CSG levels versus the control CSG levels is associated with a cancer which is regressing or in remission.

Other objects, features, advantages and aspects of the present invention will become apparent to those of skill in the art from the following description. It should be understood, however, that the following description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only. Various changes and modifications within the spirit and scope of the disclosed invention will become readily apparent to those skilled in the art from reading the following description and from reading the other parts of the present disclosure.

DESCRIPTION OF THE INVENTION

The present invention relates to diagnostic assays and methods, both quantitative and qualitative for detecting, diagnosing, monitoring, staging, and prognosticating cancers by comparing levels of CSG with those of CSG in a normal human control. What is meant by "levels of CSG" as used herein, means levels of the native protein expressed by the genes comprising the polynucleotide sequence of any of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8 or 9. In the alternative, what is meant by "levels of CSC" as used herein, means levels of the native mRNA encoded by any of the genes comprising any of the polynucleotide sequences of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8 or 9 or levels of the gene comprising any of the polynucleotide sequence of SEQ ID NO: 1, 2, 3, 4, 5, 6, 7, 8 or 9. Such levels are preferably measured in at least one of, cells, tissues and/or bodily fluids, including determination of normal and abnormal levels. Thus, for instance, a diagnostic assay in accordance with the invention for

diagnosing over-expression of any one of the CSG proteins compared to normal control bodily fluids, cells, or tissue samples may be used to diagnose the presence of cancers, including colon cancer. Any of the nine CSGs may be measured alone in the methods of the invention, or all together or any combination of the nine.

By "control" it is meant a human patient without cancer and/or non cancerous samples from the patient, also referred to herein as a normal human control; in the methods for diagnosing or monitoring for metastasis, control may also include samples from a human patient that is determined by reliable methods to have colon cancer which has not metastasized.

All the methods of the present invention may optionally include measuring the levels of other cancer markers as well as CSG. Other cancer markers, in addition to CSG, useful in the present invention will depend on the cancer being tested and are known to those of skill in the art.

Diagnostic Assays

The present invention provides methods for diagnosing the presence of colon cancer by analyzing for changes in levels of CSG in cells, tissues or bodily fluids compared with levels of CSG in cells, tissues or bodily fluids of preferably the same type from a normal human control, wherein an increase in levels of CSG in the patient versus the normal human control is associated with the presence of colon cancer. Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the patient being tested has cancer is one in which cells, tissues, or bodily fluid levels of the cancer marker, such as CSG, are at least two times higher, and most preferably are at least five times higher, than in preferably the same cells, tissues, or bodily fluid of a normal human control.

The present invention also provides a method of diagnosing metastatic colon cancer in a patient having colon

- 7 -

cancer which has not yet metastasized for the onset of metastasis. In the method of the present invention, a human cancer patient suspected of having colon cancer which may have metastasized (but which was not previously known to have metastasized) is identified. This is accomplished by a variety of means known to those of skill in the art. For example, in the case of colon cancer, patients are typically diagnosed with colon cancer following traditional detection methods.

In the present invention, determining the presence of CSG level in cells, tissues, or bodily fluid, is particularly useful for discriminating between colon cancer which has not metastasized and colon cancer which has metastasized. Existing techniques have difficulty discriminating between colon cancer which has metastasized and colon cancer which has not metastasized and proper treatment selection is often dependent upon such knowledge.

In the present invention, the cancer marker levels measured in such cells, tissues, or bodily fluid is CSG, and are compared with levels of CSG in preferably the same cells, tissue, or bodily fluid type of a normal human control. That is, if the cancer marker being observed is just CSG in serum, this level is preferably compared with the level of CSG in serum of a normal human patient. An increase in the CSG in the patient versus the normal human control is associated with colon cancer which has metastasized.

Without limiting the instant invention, typically, for a quantitative diagnostic assay a positive result indicating the cancer in the patient being tested or monitored has metastasized is one in which cells, tissues, or bodily fluid levels of the cancer marker, such as CSG, are at least two times higher, and most preferable are at least five times higher, than in preferably the same cells, tissues, or bodily fluid of a normal patient.

Normal human control as used herein includes a human patient without cancer and/or non cancerous samples from the patient; in the methods for diagnosing or monitoring for metastasis, normal human control may also include samples from a human patient that is determined by reliable methods to have colon cancer which has not metastasized.

Staging

The invention also provides a method of staging colon cancer in a human patient.

The method comprises identifying a human patient having such cancer; analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG. Then, the method compares CSG levels in such cells, tissues, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which is progressing and a decrease in the levels of CSG is associated with a cancer which is regressing or in remission.

Monitoring

Further provided is a method of monitoring colon cancer in a human having such cancer for the onset of metastasis. The method comprises identifying a human patient having such cancer that is not known to have metastasized; periodically analyzing a sample of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissue, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which has metastasized.

Further provided by this inventions is a method of monitoring the change in stage of colon cancer in a human having such cancer. The method comprises identifying a human patient having such cancer; periodically analyzing a sample

of cells, tissues, or bodily fluid from such patient for CSG; comparing the CSG levels in such cells, tissue, or bodily fluid with levels of CSG in preferably the same cells, tissues, or bodily fluid type of a normal human control sample, wherein an increase in CSG levels in the patient versus the normal human control is associated with a cancer which is progressing in stage and a decrease in the levels of CSG is associated with a cancer which is regressing in stage or in remission.

Monitoring such patient for onset of metastasis is periodic and preferably done on a quarterly basis. However, this may be more or less frequent depending on the cancer, the particular patient, and the stage of the cancer.

Assay Techniques

Assay techniques that can be used to determine levels of gene expression, such as CSG of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, reverse transcriptase PCR (RT-PCR) assays, immunohistochemistry assays, *in situ* hybridization assays, competitive-binding assays, Western Blot analyses and ELISA assays. Among these, ELISAs are frequently preferred to diagnose a gene's expressed protein in biological fluids. An ELISA assay initially comprises preparing an antibody, if not readily available from a commercial source, specific to CSG, preferably a monoclonal antibody. In addition a reporter antibody generally is prepared which binds specifically to CSG. The reporter antibody is attached to a detectable reagent such as radioactive, fluorescent or enzymatic reagent, for example horseradish peroxidase enzyme or alkaline phosphatase.

To carry out the ELISA, antibody specific to CSG is incubated on a solid support, e.g., a polystyrene dish, that binds the antibody. Any free protein binding sites on the dish are then covered by incubating with a non-specific

protein such as bovine serum albumin. Next, the sample to be analyzed is incubated in the dish, during which time CSG binds to the specific antibody attached to the polystyrene dish. Unbound sample is washed out with buffer. A reporter antibody specifically directed to CSG and linked to horseradish peroxidase is placed in the dish resulting in binding of the reporter antibody to any monoclonal antibody bound to CSG. Unattached reporter antibody is then washed out. Reagents for peroxidase activity, including a colorimetric substrate are then added to the dish. Immobilized peroxidase, linked to CSG antibodies, produces a colored reaction product. The amount of color developed in a given time period is proportional to the amount of CSG protein present in the sample. Quantitative results typically are obtained by reference to a standard curve.

A competition assay may be employed wherein antibodies specific to CSG attached to a solid support and labeled CSG and a sample derived from the host are passed over the solid support and the amount of label detected attached to the solid support can be correlated to a quantity of CSG in the sample. Nucleic acid methods may be used to detect CSG mRNA as a marker for colon cancer. Polymerase chain reaction (PCR) and other nucleic acid methods, such as ligase chain reaction (LCR) and nucleic acid sequence based amplification (NASABA), can be used to detect malignant cells for diagnosis and monitoring of various malignancies. For example, reverse-transcriptase PCR (RT-PCR) is a powerful technique which can be used to detect the presence of a specific mRNA population in a complex mixture of thousands of other mRNA species. In RT-PCR, an mRNA species is first reverse transcribed to complementary DNA (cDNA) with use of the enzyme reverse transcriptase; the cDNA is then amplified as in a standard PCR reaction. RT-PCR can thus reveal by amplification the presence of a single species of mRNA. Accordingly, if the mRNA is highly specific for the cell that produces it, RT-PCR

- 11 -

can be used to identify the presence of a specific type of cell.

Hybridization to clones or oligonucleotides arrayed on a solid support (i.e., gridding) can be used to both detect the expression of and quantitate the level of expression of that gene. In this approach, a cDNA encoding the CSG gene is fixed to a substrate. The substrate may be of any suitable type including but not limited to glass, nitrocellulose, nylon or plastic. At least a portion of the DNA encoding the CSG gene is attached to the substrate and then incubated with the analyte, which may be RNA or a complementary DNA (cDNA) copy of the RNA, isolated from the tissue of interest.

Hybridization between the substrate bound DNA and the analyte can be detected and quantitated by several means including but not limited to radioactive labeling or fluorescence labeling of the analyte or a secondary molecule designed to detect the hybrid. Quantitation of the level of gene expression can be done by comparison of the intensity of the signal from the analyte compared with that determined from known standards. The standards can be obtained by *in vitro* transcription of the target gene, quantitating the yield, and then using that material to generate a standard curve.

The above tests can be carried out on samples derived from a variety of patients' cells, bodily fluids and/or tissue extracts (homogenates or solubilized tissue) such as from tissue biopsy and autopsy material. Bodily fluids useful in the present invention include blood, urine, saliva, or any other bodily secretion or derivative thereof. Blood can include whole blood, plasma, serum, or any derivative of blood.

EXAMPLES

The present invention is further described by the following examples. These examples are provided solely to illustrate the invention by reference to specific embodiments.

- 12 -

These exemplifications, while illustrating certain specific aspects of the invention, do not portray the limitations or circumscribe the scope of the disclosed invention.

- 13 -

Example 1: CSGs

Searches were carried out and CSGs identified using the following Search Tools as part of the LIFESEQ® database available from Incyte Pharmaceuticals, Palo Alto, CA:

1. Library Comparison (compares one library to one other library) allows the identification of clones expressed in tumor and absent or expressed at a lower level in normal tissue.

2. Subsetting is similar to library comparison but allows the identification of clones expressed in a pool of libraries and absent or expressed at a lower level in a second pool of libraries.

3. Transcript Imaging lists all of the clones in a single library or a pool of libraries based on abundance. Individual clones can then be examined using Electronic Northernblots to determine the tissue sources of their component ESTs.

4. Protein Function: Incyte has identified subsets of ESTs with a potential protein function based on homologies to known proteins. Some examples in this database include Transcription Factors and Proteases. We identified some leads by searching in this database for clones whose component ESTs showed disease specificity.

Electronic subtractions, transcript imaging and protein function searches were used to identify clones, whose component ESTs were exclusively or more frequently found in libraries from specific tumors. Individual candidate clones were examined in detail by checking where each EST originated.

Table 1: CSGs

SEQ ID	Clone ID #	Gene ID #	
NO:			
1	238330	242807	Transcript Imaging
2	1285234	239588	Subsetting
3	1341701	29634	Transcript Imaging
4	816257	233421	Subsetting
5	775133	245080	Subsetting
6	1335450	245811	Subsetting
7	2348122	233711	Transcript Imaging
8	3228674	230273	Subsetting
9	1632174	229022	Transcript Imaging

The following example was carried out using standard techniques, which are well known and routine to those of skill in the art, except where otherwise described in detail. Routine molecular biology techniques of the following example can be carried out as described in standard laboratory manuals, such as Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2nd Ed.; Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989).

Example 2: Relative Quantitation of CSG Gene Expression

Real-Time quantitative PCR with fluorescent Taqman probes is a quantitation detection system utilizing the 5'-3' nuclease activity of Taq DNA polymerase. The method uses an internal fluorescent oligonucleotide probe (Taqman) labeled with a 5' reporter dye and a downstream, 3' quencher dye. During PCR, the 5'-3' nuclease activity of Taq DNA polymerase releases the reporter, whose fluorescence can then be detected by the laser detector of the Model 7700 Sequence Detection System (PE Applied Biosystems, Foster City, CA, USA).

- 15 -

Amplification of an endogenous control is used to standardize the amount of sample RNA added to the reaction and normalize for Reverse Transcriptase (RT) efficiency. Either cyclophilin, glyceraldehyde-3-phosphate dehydrogenase (GAPDH) or 18S ribosomal RNA (rRNA) is used as this endogenous control. To calculate relative quantitation between all the samples studied, the target RNA levels for one sample are used as the basis for comparative results (calibrator). Quantitation relative to the "calibrator" can be obtained using the standard curve method or the comparative method (User Bulletin #2: ABI PRISM 7700 Sequence Detection System).

To evaluate the tissue distribution, and the level of CSGs in normal and tumor tissue, total RNA was extracted from normal tissues, tumor tissues, and from tumors and the corresponding matched normal tissues. Subsequently, first strand cDNA was prepared with reverse transcriptase and the polymerase chain reaction was done using primers and Tagman probe specific to the CSG. The results were analyzed using the ABI PRISM 7700 Sequence Detector. The absolute numbers are relative levels of expression of the CSG compared to the calibrator.

Comparative Examples

Similar mRNA expression analysis for genes coding for the diagnostic markers PSA (Prostate Specific Antigen) and PLA2 (Phospholipase A2) was performed for comparison. PSA is currently the only cancer screening marker available in clinical laboratories. When the panel of normal pooled tissues was analyzed, PSA was expressed at very high levels in prostate, with a very low expression in breast and testis. After analysis of more than 55 matching samples from 14 different tissues, the data corroborated the tissue specificity seen with normal tissue samples. PSA expression was compared in cancer and normal adjacent tissue for 12 matching samples of prostate tissue. The relative levels of PSA were higher in 10 cancer samples (83%). Clinical data

- 16 -

recently obtained support the utilization of PLA2 as a staging marker for late stages of prostate cancer. mRNA expression data described herein showed overexpression of the mRNA in 8 out of the 12 prostate matching samples analyzed (66%). PLA2 had high levels of mRNA expression in small intestine, prostate, liver, and pancreas.

Measurement of SEQ ID NO:3; Clone ID 1341701; Gene ID 29634 (Cln106)

Absolute numbers are depicted in Table 2 as relative levels of expression of Cln106 (SEQ ID NO:3) in 12 normal different tissues. All the values are compared to normal testis (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 2: Relative levels of Cln106 Expression in Pooled Samples

Tissue	NORMAL
Colon-Ascending	110
Endometrium	0
Kidney	0
Liver	0
Ovary	0
Pancreas	0
Prostate	16
Small Intestine	0
Spleen	0
Stomach	0
Testis	1
Uterus	0

The relative levels of expression in Table 2 show for the CSG Cln106 (SEQ ID NO:3), mRNA expression is more than 6 fold higher in the pool of normal ascending colon (110) compared with prostate (16). Testis, the calibrator, with a relative expression level of 1, is the only other tissue expressing the mRNA for Cln106 (SEQ ID NO:3). These results demonstrate that mRNA expression of this CSG is highly specific for colon.

- 17 -

The absolute numbers in Table 2 were obtained analyzing pools of samples of a particular tissue from different individuals. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 3.

The absolute numbers in Table 3 are relative levels of expression of Cln106 (SEQ ID NO:3) in 57 pairs of matching samples. All the values are compared to normal testis (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 3: Relative levels of Cln106 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matching Normal Adjacent
Sto AC93	Stomach 1	4	96
Sto AC99	Stomach 2	0.4	0.5
Sml 21XA	Small Intestine 1	0	0
Sml H89	Small Intestine 2	0.93	1.28
Cln B56	Colon-Cecum(A) 1	317	101
Cln AS45	Colon-Ascending(A) 2	316.3	146.5
Cln CM67	Colon-Cecum(B) 3	481.0	217.5
Cln AS67	Colon-Ascending(B) 4	858.1	220.6
Cln AS43	Colon-Ascending(C) 5	1370	98
Cln AS46	Colon-Ascending(C) 6	3051	375
Cln AS98	Colon-Ascending(C) 7	26	42
Cln AS89	Colon-Ascending(D) 8	524.6	11.0
Cln TX01	Colon-Transverse(B) 9	2886.3	1992.0
Cln TX89	Colon-Transverse(B) 10	146.0	35.9
Cln TX67	Colon-Transverse(C) 11	2.9	421.7
Cln MT38	Colon-Splenic Flexure(M) 12	1681	187

Cln SG89	Colon-Sigmoid(B)13	1063.8	31.1
Cln SG67	Colon-Sigmoid(C)14	8.5	9.4
Cln SG33	Colon-Sigmoid(C)15	264	549
Cln SG45	Colon-Sigmoid(D)16	580.0	114.6
Cln B34	Colon-Rectosigmoid(A)17	97	244
Cln CXGA	Colon-Rectum(A)18	45.1	273.4
Cln RC67	Colon-Rectum(B)19	2.7	20.0
Cln C9XR	Colon-Rectosigmoid(C)20	609	460
Cln RS45	Colon-Rectosigmoid(C)21	472.8	144.0
Cln RC01	Colon-Rectum(C)22	568	129
Cln RC89	Colon-Rectum(D)23	4.6	322.91
Bld 46XK	Bladder 1	0.2	0
Bld 66X	Bladder 2	1	1
Bld 32XK	Bladder 3	0.0	0.0
Kid 126XD	Kidney 1	0	0
Kid 12XD	Kidney 2	0	0
Kid 5XD	Kidney 3	0.0	1.0
Kid 6XD	Kidney 4	0.0	0.0
Kid 106XD	Kidney 5	0.4	0.0
Liv 42X	Liver 1	0.0	0.0
Liv 15XA	Liver 2	0.0	0.0
Liv 94XA	Liver 3	0.0	0.0
Lng AC69	Lung 1	2	0
Lng BR94	Lung 2	0	0
Lng 47XQ	Lung 3	0	0
Mam 59X	Mammary Gland 1	0	0
Mam B011X	Mammary Gland 2	0	0
Mam A06X	Mammary Gland 3	0	0
Ovr 103X	Ovary 1	0.04	2.08
Ovr 130X	Ovary 2	0.1	2.76

- 19 -

Pan 71XL	Pancreas 1	4.08	0.1
Pan 82XP	Pancreas 2	0	0
Pro 12B	Prostate 1	0.3	0
Pro 23B	Prostate 2	3	4
Pro 13XB	Prostate 3	2	7
Pro 34B	Prostate 4	0.54	4.01
Pro 20XB	Prostate 5	4.8	4.3
Pro 65XB	Prostate 6	0.7	1.3
Tst 39X	Testis 1	2.78	0
End 8XA	Endometrium 1	0	0.2
Utr 85XU	Uterus 1	1.26	0

0= Negative

When matching samples were analyzed, the higher levels of expression were in the colon, showing a high degree of tissue specificity for this tissue. These results confirm the tissue specificity results obtained with the panel of normal pooled samples (Table 2). Furthermore, the level of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer stage (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 3 shows overexpression of Cln106 (SEQ ID NO:3) in 15 colon cancer tissues compared with their respective normal adjacent (colon samples #1, 2, 3, 4, 5, 6, 8, 9, 10, 12, 13, 16, 20, 21, and 22). There is overexpression in the cancer tissue for 65% of the colon matching samples tested (total of 23 colon matching samples). The matching sample Pan 71XL is a secondary cancer in pancreas, the primary cancer in that individual was a duodenal cancer.

Altogether, the high level of tissue specificity, plus the mRNA overexpression in 65% of the colon matching samples

- 20 -

tested are demonstrative of CSG Cln106 (SEQ ID NO:3) being a diagnostic marker for colon cancer.

Measurement of SEQ ID NO:4; Clone ID 816257; Gene ID 406452 (Cln107)

Absolute numbers as depicted in Table 4 are relative levels of expression of CSG Cln107 (SEQ ID NO:4) in 12 normal different tissues. All the values are compared to normal small intestine (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 4: Relative levels of Cln107 Expression in Pooled Samples

Tissue	NORMAL
Colon-Ascending	3.2
Endometrium	0
Kidney	0.2
Liver	0
Ovary	0
Pancreas	0
Prostate	0.1
Small Intestine	1
Spleen	0
Stomach	0.3
Testis	0
Uterus	0

The relative levels of expression in Table 4 show that mRNA expression of the CSG Cln107 (SEQ ID NO:4) is more than 10 fold higher in the pool of normal ascending colon (3.2), five fold higher in small intestine (1), and 1.5 fold higher in stomach (0.3), compared with the next higher expressor (0.2 for kidney). Seven of the pooled tissues samples analyzed were negative and prostate showed a relative expression of 0.1 for Cln107 (SEQ ID NO:4). These results demonstrate that Cln107 mRNA expression is highly specific for colon, small intestine, and in a lower degree for stomach.

The absolute numbers in Table 4 were obtained analyzing pools of samples of a particular tissue from different

- 21 -

individuals. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 5.

The absolute numbers in Table 5 are relative levels of expression of Cln107 (SEQ ID NO:4) in 57 pairs of matching samples. All the values are compared to normal small intestine (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 5: Relative levels of Cln107 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matching Normal Adjacent
Sto AC93	Stomach 1	8.9	13.4
Sto AC99	Stomach 2	6.0	0.9
Sml 21XA	Small Intestine 1	1.07	1.42
Sml H89	Small Intestine 2	0.97	4.13
Cln B56	Colon-Cecum(A) 1	2	16
Cln AS45	Colon-Ascending(A) 2	0.7	2.1
Cln CM67	Colon-Cecum(B) 3	1.6	2.1
Cln AS67	Colon-Ascending(B) 4	1.2	6.2
Cln AS43	Colon-Ascending(C) 5	13.5	0.5
Cln AS46	Colon-Ascending(C) 6	9.7	23.6
Cln AS98	Colon-Ascending(C) 7	28.1	1.4
Cln AS89	Colon-Ascending(D) 8	0.9	3.1
Cln TX01	Colon-Transverse(B) 9	3.0	10.6
Cln TX89	Colon-Transverse(B) 10	4.5	0.6
Cln TX67	Colon-Transverse(C) 11	3.6	3.4
Cln MT38	Colon-Splenic Flexure(M) 12	4.0	2.6
Cln SG89	Colon-Sigmoid(B) 13	4.7	0.9

Cln SG67	Colon-Sigmoid(C)14	1.0	1.3
Cln SG33	Colon-Sigmoid(C)15	14.2	7.6
Cln SG45	Colon-Sigmoid(D)16	4.8	6.0
Cln B34	Colon-Rectosigmoid(A)17	3	2
Cln CXGA	Colon-Rectum(A)18	4.4	1.9
Cln RC67	Colon-Rectum(B)19	0.1	0.4
Cln C9XR	Colon-Rectosigmoid(C)20	5	3
Cln RS45	Colon-Rectosigmoid(C)21	11.4	4.6
Cln RC01	Colon-Rectum(C)22	1.8	2.3
Cln RC89	Colon-Rectum(D)23	0.1	5.35
Bld 46XK	Bladder 1	0.2	0
Bld 66X	Bladder 2	1	1
Bld 32XK	Bladder 3	0.1	0.1
Kid 126XD	Kidney 1	0	0.02
Kid 12XD	Kidney 2	0.1	0.2
Kid 5XD	Kidney 3	0.3	0.0
Kid 6XD	Kidney 4	0.1	0.1
Kid 106XD	Kidney 5	0.0	0.1
Liv 42X	Liver 1	7.9	0.002
Liv 15XA	Liver 2	0.0	0.0
Liv 94XA	Liver 3	0.0	0.0
Lng AC69	Lung 1	1.6	0.2
Lng BR94	Lung 2	0.4	0
Lng 47XQ	Lung 3	0.78	0.2
Mam 59X	Mammary Gland 1	0.05	0.3
Mam B011X	Mammary Gland 2	0.01	0.004
Mam A06X	Mammary Gland 3	0.22	0
Ovr 103X	Ovary 1	0.01	0.01
Ovr 130X	Ovary 2	0.09	0.1
Pan 71XL	Pancreas 1	2.51	2.81

- 23 -

Pan 82XP	Pancreas 2	0	0.62
Pro 12B	Prostate 1	0.3	0.1
Pro 23B	Prostate 2	0.3	0.2
Pro 13XB	Prostate 3	0	0
Pro 34B	Prostate 4	0.04	0.22
Pro 20XB	Prostate 5	0.4	0.1
Pro 65XB	Prostate 6	0.0	0.1
Tst 39X	Testis 1	0.02	0.01
End 8XA	Endometrium 1	0.01	0.5
Utr 85XU	Uterus 1	0.03	0

0= Negative

When matching samples were analyzed, the higher levels of expression were in colon, stomach, and small intestine, showing a high degree of tissue specificity for colon tissues. These results confirm the tissue specificity results obtained with normal pooled samples (Table 4). Furthermore, the level of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer stage (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 5 shows overexpression of Cln107 (SEQ ID NO:4) in 11 colon cancer tissues compared with their respective normal adjacent (colon samples #5, 7, 10, 11, 12, 13, 15, 17, 18, 20, and 21). There is overexpression in the cancer tissue for 48% of the colon matching samples tested (total of 23 colon matching samples). The matching sample Pan 71XL is a secondary cancer in pancreas, the primary cancer in that individual was a duodenal cancer.

Altogether, the high level of tissue specificity, plus the mRNA overexpression in almost half of the colon, stomach, and small intestine matching samples tested are demonstrative

- 24 -

of CSG Cln107 (SEQ ID NO:4) being a diagnostic marker for colon cancer.

Measurement of SEQ ID NO:5; Clone ID 775133; Gene ID 24508 (Cln108)

The absolute numbers shown in Table 6 are relative levels of expression of CSG Cln108 (SEQ ID NO:5) in 12 normal different tissues. All the values are compared to normal small intestine (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 6: Relative levels of Cln108 Expression in Pooled Samples

Tissue	NORMAL
Colon-Ascending	2846.5
Endometrium	1
Kidney	5.5
Liver	18.7
Ovary	3.4
Pancreas	198.1
Prostate	1024
Small Intestine	810.8
Spleen	32.2
Stomach	9981.2
Testis	0
Uterus	294.1

The relative levels of expression in Table 6 show that mRNA expression of CSG Cln108 (SEQ ID NO:5) is more than 10 fold higher in the pool of normal ascending colon (2846.5) and almost ten fold higher in stomach (9981.2), compared to the expression level in any other tissue analyzed. These results demonstrate that mRNA expression of this CSG is also highly specific for colon and stomach.

The absolute numbers in Table 6 were obtained analyzing pools of samples of a particular tissue from different individuals. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 7.

The absolute numbers depicted in Table 7 are relative levels of expression of Cln108 (SEQ ID NO:5) in 57 pairs of matching samples. All the values are compared to normal small intestine (calibrator). A matching pair is formed by mRNA from the cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 7: Relative levels of Cln108 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matching Normal Adjacent
Sto AC93	Stomach 1	28696	34842
Sto AC99	Stomach 2	21523	30862
Sml 21XA	Small Intestine 1	2944	964.4
Sml H89	Small Intestine 2	244.5	3513.2
Cln B56	Colon-Cecum(A) 1	27242	24637
Cln AS45	Colon-Ascending(A) 2	5827.0	8771.0
Cln CM67	Colon-Cecum(B) 3	4251.0	4684.0
Cln AS67	Colon-Ascending(B) 4	564.0	1949.0
Cln AS43	Colon-Ascending(C) 5	50310	10949
Cln AS46	Colon-Ascending(C) 6	246044	120073
Cln AS98	Colon-Ascending(C) 7	40442	17482
Cln AS89	Colon-Ascending(D) 8	5730.0	1581.0
Cln TX01	Colon-Transverse(B) 9	22281.0	114784.0
Cln TX89	Colon-Transverse(B) 10	11026.0	1639.0
Cln TX67	Colon-Transverse(C) 11	17004.0	11654.0
Cln MT38	Colon-Splenic Flexure(M) 12	77589	31620
Cln SG89	Colon-Sigmoid(B) 13	140339.0	49617.0
Cln SG67	Colon-Sigmoid(C) 14	4951.0	7905.0
Cln SG33	Colon-Sigmoid(C) 15	60875	120490

Cln SG45	Colon-Sigmoid(D)16	30437.0	47267.0
Cln B34	Colon-Rectosigmoid(A)17	5848	5861
Cln CXGA	Colon-Rectum(A)18	13877.0	9787.0
Cln RC67	Colon-Rectum(B)19	1703.0	26589.0
Cln C9XR	Colon-Rectosigmoid(C)20	2458	19071
Cln RS45	Colon-Rectosigmoid(C)21	95523	61939
Cln RC01	Colon-Rectum(C)22	98891.0	80047.0
Cln RC89	Colon-Rectum(D)23	17.0	1775
Bld 46XK	Bladder 1	0	8
Bld 66X	Bladder 2	397	44
Bld 32XK	Bladder 3	0.0	16.0
Kid 126XD	Kidney 1	32	22
Kid 12XD	Kidney 2	6	0
Kid 106XD	Kidney 3	4.0	33.0
Liv 42X	Liver 1	4783	0
Liv 15XA	Liver 2	4.0	10.0
Liv 94XA	Liver 3	159.0	21.0
Lng AC69	Lung 1	222	295
Lng BR94	Lung 2	112	0
Lng 47XQ	Lung 3	30	69
Lng AC66	Lung 4	29	137
Mam 59X	Mammary Gland 1	56	0
Mam B011X	Mammary Gland 2	54	31
Mam A06X	Mammary Gland 3	12	0
Ovr 103X	Ovary 1	37	0
Pan 71XL	Pancreas 1	13203	4163
Pan 82XP	Pancreas 2	39.1	0
Pro 12B	Prostate 1	386	88
Pro 23B	Prostate 2	250	23
Pro 13XB	Prostate 3	92	731

- 27 -

Pro 34B	Prostate 4	33.3	265.7
Pro 20XB	Prostate 5	454.6	1908.9
Pro 65XB	Prostate 6	733.5	922.0
End 8XA	Endometrium 1	5	92
Utr 85XU	Uterus 1	98.9	21.8
Utr 23XU	Uterus 2	35.3	0
Utr 135XO	Uterus 3	39.2	43.8
Utr 141XO	Uterus 4	212.1	55.9

0= Negative

When matching samples were analyzed, the higher levels of expression were in colon and stomach, showing a high degree of tissue specificity for these two tissues. These results confirm the tissue specificity results obtained with normal pooled samples (Table 6). Furthermore, the level of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer stage (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 7 shows overexpression of CSG Cln108 (SEQ ID NO:5) in 13 colon cancer tissues compared with their respective normal adjacent (colon samples #1, 5, 6, 7, 8, 9, 10, 11, 12, 13, 18, 21, and 22). There is overexpression in the cancer tissue for 56% of the colon matching samples tested (total of 23 colon matching samples). The matching sample Pan 71XL is a secondary cancer in pancreas, the primary cancer in that individual was a duodenal cancer.

Altogether, the high level of tissue specificity, plus the mRNA overexpression in more than half of the colon, stomach, and small intestine matching samples tested are demonstrative of this CSG, Cln108 (SEQ ID NO:5), also being a diagnostic marker for colon cancer.

**Measurement of SEQ ID NO:7; Clone ID 2348122; Gene ID 23371
(Cln109)**

The absolute numbers depicted in Table 8 are relative levels of expression of CSG Cln109 (SEQ ID NO:7) in 12 normal different tissues. All the values are compared to normal ovary (calibrator). These RNA samples are commercially available pools, originated by pooling samples of a particular tissue from different individuals.

Table 8: Relative levels of Cln109 Expression in Pooled Samples

Tissue	NORMAL
Colon-Ascending	28.8
Endometrium	0.45
Kidney	0.41
Liver	0.72
Ovary	0.07
Pancreas	82.8
Prostate	124.3
Small Intestine	626.4
Spleen	1.2
Stomach	12.05
Testis	1.51
Uterus	52.99

The relative levels of expression in Table 8 show that mRNA expression of CSG Cln109 (SEQ ID NO:7), is more than 5 fold higher in the pool of normal small intestine (626.4) compared to the expression level in any other tissue analyzed. These results demonstrate that Cln109 (SEQ ID NO:7) mRNA expression is highly specific for small intestine

The absolute numbers in Table 8 were obtained analyzing pools of samples of a particular tissue from different individuals. They can not be compared to the absolute numbers originated from RNA obtained from tissue samples of a single individual in Table 9.

The absolute numbers depicted in Table 9 are relative levels of expression of Cln109 (SEQ ID NO:7) in 53 pairs of matching samples. All the values are compared to normal ovary (calibrator). A matching pair is formed by mRNA from the

cancer sample for a particular tissue and mRNA from the normal adjacent sample for that same tissue from the same individual.

Table 9: Relative levels of Cln109 Expression in Individual Samples

Sample ID	Tissue	Cancer	Matching Normal Adjacent
Sto AC93	Stomach 1	2574	1310
Sto AC99	Stomach 2	4153	5
Sml 21XA	Small Intestine 1	2667	13663.8
Sml H89	Small Intestine 2	57.8	904.29
Cln B56	Colon-Cecum(A) 1	6794	299
Cln AS45	Colon-Ascending(A) 2	814.6	105.8
Cln CM67	Colon-Cecum(B) 3	294.6	36.1
Cln AS67	Colon-Ascending(B) 4	2.2	26.3
Cln AS43	Colon-Ascending(C) 5	111	377
Cln AS46	Colon-Ascending(C) 6	1180	352
Cln AS98	Colon-Ascending(C) 7	1075	92
Cln AS89	Colon-Ascending(D) 8	14022.7	87.5
Cln TX01	Colon-Transverse(B) 9	1027.6	282.1
Cln TX89	Colon-Transverse(B) 10	2.5	23.7
Cln TX67	Colon-Transverse(C) 11	0.1	72.3
Cln MT38	Colon-Splenic Flexure(M) 12	372	88
Cln SG89	Colon-Sigmoid(B) 13	179.2	33.4
Cln SG67	Colon-Sigmoid(C) 14	85.0	94.7
Cln SG33	Colon-Sigmoid(C) 15	5461	377
Cln SG45	Colon-Sigmoid(D) 16	762.7	15.9
Cln B34	Colon-Rectosigmoid(A) 17	460	1
Cln RC67	Colon-Rectum(B) 18	64.5	136.2
Cln C9XR	Colon-Rectosigmoid(C) 19	441	34
Cln RS45	Colon-Rectosigmoid(C) 20	1931	195

Cln RC01	Colon-Rectum(C)21	72.8	19.1
Cln RC89	Colon-Rectum(D)22	4.8	90.2
Bld 46XK	Bladder 1	4	3
Bld 66X	Bladder 2	1	0
Bld 32XK	Bladder 3	0.1	307.6
Kid 126XD	Kidney 1	0	2
Kid 12XD	Kidney 2	3	16
Kid 5XD	Kidney 3	0.0	0.3
Kid 6XD	Kidney 4	18.5	1.2
Liv 42X	Liver 1	21	0.03
Liv 15XA	Liver 2	0.5	0.4
Liv 94XA	Liver 3	0.4	0.0
Lng AC69	Lung 1	0.1	0
Lng BR94	Lung 2	3	0
Lng 60XL	Lung 3	0.1	0
Mam 59X	Mammary Gland 1	0	4
Mam B011X	Mammary Gland 2	8	13
Mam A06X	Mammary Gland 3	4.7	9.6
Pan 71XL	Pancreas 1	8902.5	1428.2
Pan 82XP	Pancreas 2	0.2	9.3
Pro 12B	Prostate 1	9	20
Pro 23B	Prostate 2	191	88
Pro 13XB	Prostate 3	12	460
Pro 34B	Prostate 4	3.2	80.4
Tst 39X	Testis 1	29.9	0
End 8XA	Endometrium 1	0.3	21
Utr 85XU	Uterus 1	244.7	592.2
Ovr 63A	Ovary 1	11.4	0
Ovr A1C	Ovary 2	68.4	0

0= Negative

- 31 -

When matching samples were analyzed, the higher levels of expression were in small intestine, colon and stomach, showing a high degree of tissue specificity for these three colon tissues. These results confirm the tissue specificity results obtained with normal pooled samples for small intestine (Table 8). Furthermore, the level of mRNA expression in cancer samples and the isogenic normal adjacent tissue from the same individual were compared. This comparison provides an indication of specificity for the cancer stage (e.g. higher levels of mRNA expression in the cancer sample compared to the normal adjacent). Table 9 shows overexpression of CSG, Cln109 (SEQ ID NO:7) in 15 colon cancer tissues compared with their respective normal adjacent (colon samples #1, 2, 3, 6, 7, 8, 9, 12, 13, 15, 16, 17, 19, 20, and 21). There is overexpression in the cancer tissue for 68% of the colon matching samples tested (total of 22 colon matching samples). The matching sample Pan 71XL is a secondary cancer in pancreas, the primary cancer in that individual was a duodenal cancer.

Altogether, the high level of tissue specificity, plus the mRNA overexpression in more than half of the colon, stomach, and small intestine matching samples tested are demonstrative of CSG Cln109 (SEQ ID NO:7) being a diagnostic marker for colon cancer. The amino acid sequence encoded by the open reading frame of Cln109 is depicted in SEQ ID NO:10.

- 32 -

What is claimed is:

1. A method for diagnosing the presence of colon cancer in a patient comprising:

(a) measuring levels of CSG in a sample of cells, tissue or bodily fluid obtained from the patient; and

(b) comparing the measured levels of CSG with levels of CSG in a sample of cells, tissue or bodily fluid obtained from a control, wherein an increase in measured levels of CSG in the patient versus the CSG levels in the control is associated with the presence of colon cancer.

2. A method of diagnosing metastatic colon cancer in a patient comprising:

(a) measuring levels of CSG in a sample of cells, tissue, or bodily fluid obtained from the patient; and

(b) comparing the measured levels of CSG with levels of CSG in a sample of cells, tissue, or bodily fluid obtained from a control, wherein an increase in measured CSG levels in the patient versus the CSG levels in the control is associated with a cancer which has metastasized.

3. A method of staging colon cancer in a patient comprising:

(a) identifying a patient suffering from colon cancer;

(b) measuring levels of CSG in a sample of cells, tissue, or bodily fluid obtained from the patient; and

(c) comparing the measured levels of CSG with levels of CSG in a sample of cells, tissue, or bodily fluid obtained from a control, wherein an increase in the measured levels of CSG versus the levels of CSG in the control is associated with a cancer which is progressing and a decrease in the measured levels of CSG versus the levels of CSG in the control is associated with a cancer which is regressing or in remission.

- 33 -

4. A method of monitoring colon cancer in a patient for the onset of metastasis comprising:

(a) identifying a patient having colon cancer that is not known to have metastasized;

(b) periodically measuring CSG levels in samples of cells, tissue, or bodily fluid obtained from the patient; and

(c) comparing the periodically measured levels of CSG with levels of CSG in cells, tissue, or bodily fluid obtained from a control, wherein an increase in any one of the periodically measured levels of CSG in the patient versus the levels of CSG in the control is associated with a cancer which has metastasized.

5. A method of monitoring changes in a stage of colon cancer in a patient comprising:

(a) identifying a patient having colon cancer;

(b) periodically measuring levels of CSG in samples of cells, tissue, or bodily fluid obtained from the patient; and

(c) comparing the measured levels of CSG with levels of CSG in a sample of the same cells, tissue, or bodily fluid of a control, wherein an increase in any one of the periodically measured levels of CSG versus levels of CSG in the control is associated with a cancer which is progressing in stage and a decrease in any one of the periodically measured levels of CSG versus the levels of CSG in the control is associated with a cancer which is regressing in stage or in remission.

6. The method of claim 1, 2, 3, 4 or 5 wherein the CSG comprises SEQ ID NO:3, 4, 5 or 7.

SEQUENCE LISTING

<110> Macina, Roberto A.
 Yang, Fei
 Sun, Yongming

<120> A Novel Method of Diagnosing, Monitoring and Staging
 Colon Cancers

<130> DEX-0035

<140>

<141>

<150> 60/086,266

<151> 1998-05-21

<160> 10

<170> PatentIn Ver. 2.0

<210> 1

<211> 487

<212> DNA

<213> Homo sapiens

<400> 1

```
tctgcatctg gccctcccag tgcacctgtt caatcccagc ycctccctga cctgtacaaa 60
tacacctgag gaccggctcg agcccagact tcctgcccct gctctgcact ctcaggtatt 120
ccctgctctt actccaaaaa gatggacca ggtccgaagg ggcactgcca ctgtgggggg 180
catggccatc ctccagggtca ctgcggggcga acccctggcc atggcccagg gccctgcggg 240
ccaccccctg gccatggccc agggccctgc gggcaacccc ctggccatgg ccagggccc 300
tgcgggcctc ccctgggcca tggcccaggt caccaccccc ctggtccaca tcaactgagga 360
agtagaagaa aacaggacac aagatggcaa gcctgagaga attgcccagc tgacctggaa 420
tgaggcctaa accacaatct tctcttcta ataaacagcc tcctagagge cacattctat 480
tctttaa                                     487
```

<210> 2

<211> 739

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (693)

<220>

<221> unsure

<222> (698)..(699)

<220>

<221> unsure

<222> (703)..(705)

<220>

<221> unsure

<222> (708)

<220>

<221> unsure

<222> (710)..(716)

<220>

<221> unsure

<222> (718)

<220>

<221> unsure

<222> (723)..(726)

<220>

<221> unsure

<222> (728)

<220>

<221> unsure

<222> (732)

<220>

<221> unsure

<222> (737)

<400> 2

```

tctgaaactg tcagttccac cagcaactgt tggatactgg taagtttcca gggggctgct 60
ttgcatctga aactgtcagc cccagaatgt tgacagtgcg tctctagcc cttctctgtg 120
cctcagcctc tggcaatgcc attcaggcca ggtcttctc ctatagtggg gagtatggaa 180
gtgggtgggtg aaagcgattc tctcattctg gcaaccagtt ggacggcccc atcacgccc 240
tccgggtccg agtcaacaca tactacatcg taggtcttca ggtgcgctat ggcaaggtgt 300
ggagcgacta tgtgggtggt cgcaacggag acctggagga gatctttctg caccctgggg 360
aatcagtgat ccaggtttct gggaagtaca agtggtagct gaagaagctg gtatttgtga 420
cagacaaggg ccgctatctg tcttttggga aagacagtgg cacaagtttc aatgcctccc 480
ccttgcaccc caacacogtg ctccgcttca tcagtggcgg gtctggttct ctcatcgatg 540
ccattggcct ccaactgggat gtttaccoca ctagtgcag cagatgctga gcctcctctc 600
cttggcaggg gcactgtgat gaggagtaag aactccctta tcaactaaccc ccattccaaat 660
ggctcaataa aaaaatatgg ttaaggctaa aaaaaaannn gannnaanan nnnnnntnca 720
aannnnantt cncctgnta 739

```

<210> 3
 <211> 428
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> (391)

<400> 3
 aattgtccgg ggtcaaacag aggagagcat gaatgagagt catcctcgca agtgtgcaga 60
 gtcttttgag atgtgggatg atcgtgactc ccactgtagg cgccttaagt ttgaaggcca 120
 tccccctgag tcttggaagt ggatccttgc accggtcatt ctttatatct gtgaaaggat 180
 cctccgggtt tcccgctccc agcagaaggt tgtgattacc aagggtgtta tgcacccatc 240
 caaagttttg gaattgcaga tgaacaagcg tggcttcagc atggaagtgg ggcagtatat 300
 ctttgttaat tgcacctcaa tctctctcct gggaatggca tcccttactt ttgacctctg 360
 ctccagagga agatttcttc ttcattcata tncgagcagc aggggacttg acagaaaatc 420
 tataagggg 428

<210> 4
 <211> 1347
 <212> DNA
 <213> Homo sapiens

<400> 4
 ggaaaacccc tgagcacaaa gcaagaggca tcgaagcccc ctcggggatg cccgcaagcc 60
 aacaggggtg tctgtcgggt ggagtacttc cgcctgcgtc ctctgcggtt cagggcccca 120
 gacgagcccc agcaggccca agtccccat gtctggggtt gggaggtggc tggggccctt 180
 gcactgaggg tgcagaagtc ccagtcactt gatctgctgg aaaggagag ggagagtgtc 240
 ctgcgcgggg agcaagaggt ggcagaggag cggagaaatg ctctcttccc agaggtcttc 300
 tccccaacgc cagatgagaa ctctgaccag aactccagga gctcctccca gccatccggc 360
 atcacgggca gttactcggg gtctgagtct ccttcttcca gccccatcca cctacactca 420
 aacgtggcgt ggacagtggg agatccagtg gacagtgtc ctcccgggca gagaaagaag 480
 gagcaatggt acgttggcat caaccctcg gacggtatca actcagagg cctggaagcc 540
 atacgggtga cccgtcaca gaacgccatg gcagagcgtt gggaatccc catctacgcc 600
 agtgaggagg atgactgagc ctcgggatgg ggcgcccacc cctgcccctg cctgaccct 660
 cgtgggaact gccaaagcca tcgccaaagc cccaccctag gaaatgggtc ctagggtccag 720
 gatccaagaa ccacagctca tctgccaa ca tcccaccat gggcacattt gggactgttg 780
 ggtttttcgt ttccgtttct atcttcttt agaaatgtt ctgccttttg ggtctaaaag 840
 ttttggggat gaaatgggga cccctgctga ttctttctgc ttctaagact ttgccaaatg 900
 ccttgggtct aagaaagaaa gagaccgct cctccacttt caggtgtaat ttgcttccgc 960
 tagtctgagg gcagagggac cgggtcaaaga ggggtggcaca gatcgagca ccttgagggg 1020
 ctgcgggtct gagggaggag acaactcagct cctccctctg agaagtccca agctgagagg 1080
 ggagacctgc ccttttccaa ccttgggaaa ccatccagtc tgaggaggga ggccaaactc 1140
 ccagtgtctg ggttccctgt gcagccctca aacccttcac cttggtgcac ccagccacac 1200
 ctggtggaca caaagctctc acatcgatag gatcccatga ggatggtccc cttcacctgg 1260
 gagaaaagtg acccagttta ggagctggag ggggtcttt gtccccacc cccaaactgc 1320

cctgaaataa acctggagtg agctgcc

1347

<210> 5

<211> 1249

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (1034)..(1046)

<400> 5

```

ggcagagcct ggcagggcca ggagcagctg gccactggc ggcccgcaac actccgtctc 60
accctctggg cccactggat ctagaaggag gccgactgtg aggccactac ccctccagca 120
actgggaggt gggactgtca gaagctggcc caggggtgtg gtcagctggg tcagggacct 180
aacggcacct ggcagggaac acctgcctt ctccatcgaa gcaggggaag tgggagcctc 240
gagccctcgg gtggaagctg accccaagcc acccttcacc tggacaggat gagagtgtca 300
ggtgtgtctt gcctcctggc cctcatctt gccatagtoa cgacatggat gtttatttga 360
agctacatga gcttcagcat gaaaaccatc cgtctgccac gctggctggc ctgccacc 420
aaggagatcc aggttaaaaa gtacaagtgt ggctcatca agccctgcc agccaactac 480
tttgcgttta aaatctgcag tggggccgcc aacgtcgtg gccctactat gtgctttgaa 540
gaccgcatga tcatgagtc tgtgaaaaac aatgtgggca gaggcctaaa catcgccctg 600
gtgaatggaa ccacgggagc tgtgctggga cagaagtcac ttgacatgta ctctggagat 660
gttatgcacc tagtgaaatt ccttaaagaa attccggggg gtgcaactgt gctgggtggc 720
tcctacgacg atccagggac caaatgaac gatgaaagca ggaaactctt ctctgacttg 780
gggagttcct acgcaaaaaca actgggcttc cgggacagct gggctctcat aggagccaaa 840
gacctcaggg gtaaaagccc ctttgagcag ttcttaaaga acagcccaga cacaaacaaa 900
tacgagggat ggcagagct gctggagatg gagggtgca tgcccccga gccattttag 960
ggtggtgtg gctcttctc agccaggggc ctgaagaagc tctgctga cttaggagtc 1020
agagcccggc agnnnnnnnn nnnnnnnnn nnnnnntgt gcgtggaagg tgcagcaggt 1080
ccttgcaagc tgtgtcgcgc ctctctctc cggaaacaga accctcccac agcacatcct 1140
acccggaaga ccagcctcag agggctctt tggaaaccag tgtctgtgga gagaatgggg 1200
tgctttcgtc aggaactgt gacggctgtt cctgaqqaag gacaaactg 1249

```

<210> 6

<211> 1220

<212> DNA

<213> Homo sapiens

<400> 6

```

gctttctgca cctcattcca catcaggagc gtttttggag aaagctgca tctgttgagc 60
tccagggcgc agtggaggga gggagtgaag gagctctctg taccgaagga aagtgcagct 120
gagactcaga caagattaca atgaaccaac tcagcttctt gctgtttctc atagegacca 180
ccagaggatg gagtacagat gaggctaata ctacttcaa ggaatggacc tgttcttctg 240
ctccatctct gccagaagc tgcaaggaaa tcaaagacga atgtctagt gcatttgatg 300
gcctgtatct tctccgcaat gagaatggtg ttatctacca gaccttctgt gacatgacct 360
ctgggggtgg cggctggacc ctggtggcca gcgtgcaoga gaatgacatg cgtgggaagt 420
gcacggtggg cgatcgctgg tccagtcagc agggcagcaa agcagctctac ccagaggggg 480

```

```

acggcaactg ggccaactac aacacctttg gatctgcaga ggccggccacg agcgatgact 540
acaagaaccc tggctactac gacatccagg ccaaggacct gggcatctgg cacgtgcccc 600
ataagtcacc catgcagcac tggagaaaca gctccctgct gaggtaccgc acggacactg 660
gcttcctcca gacactggga cataatctgt ttggcatcta ccagaaatat ccagtgaat 720
atggagaagg aaagtgttgg actgacaacg gcccggtgat cctgttgggc tatgattttg 780
ggcagcggca gaaaacagca tcttattact caccctatgg ccagcgggaa ttcactgccc 840
gatttgttca gttcagggtt ttaataacg agagagcagc caacgccttg tgtgctggaa 900
tgagggtcac cggatgtaac actgagcacc actgcatttg tggaggagga tactttccag 960
aggccagtc cagcagctgt ggagattttt ctggttttga ttgagtgga tatggaactc 1020
atgttggtta cagcagcagc cgtgagataa ctgaggcagc tgtgcttcta tctatctgt 1080
gagagttttg tggaggggaa ccagacctc tctcccaac catgagatcc caaggatgga 1140
gaacaactta ccagtagct agaattgtta tggcagaaga gaaaacaata aatcatattg 1200
actcaaaaaa aaaaaaaaag                                     1220

```

<210> 7

<211> 2796

<212> DNA

<213> Homo sapiens

<400> 7

```

cggctcgagg gacaggatga ggcccgccct ctcatctctc ctacccttc tgttcttctc 60
tggccaagct gcaggggatt tgggggatgt gggacctcca attcccagcc cgggcttcag 120
ccctttccca ggtgttgact ccagctccag ctccagctcc agctccaggt cgggctccag 180
ctccagccgc agcttaggca gcggagggtc tgtgtccag ttgttttcca atttcaccgg 240
ctccgtggat gaccgtggga cctgccagt ctctgtttcc ctgccagaca ccacctttcc 300
cgtggacaga gtggaacgct tgggaattcac agctcatgtt ctctctcaga agtttgagaa 360
agaactttcc aaagtggagg aatatgtcca attaattagt gtgtatgaaa agaaactgtt 420
aaaccttaact gtccgaattg acatcatgga gaaggatacc atttcttaca ctgaactgga 480
cttcgagctg atcaaggtag aagtgaagga gatggaaaaa ctggtcatac agctgaagga 540
gagttttggt ggaagctcag aaattgttga ccagctggag gtggagataa gaaatatgac 600
tctcttggtg gagaagcttg agacactaga caaaaacaat gtcccttgcca ttccgagaga 660
aatcgtggct ctgaagacca agctgaaaga gtgtgaggcc tctaaagatc aaaacacccc 720
tgtcgtccac cctctcccca ctccagggag ctgtgggtcat ggtgggtgtg tgaacatcag 780
caaacctgtc gtggttcagc tcaactggag aggggtttct tatctatatg gtgcttgggg 840
tagggattac tctcccagc atccaaacaa aggaactgtat tgggtggcgc cattgaatac 900
agatgggaga ctgttggaat attatagact gtacaacaca ctggatgatt tgcatttga 960
tataaatgct cgagagttgc ggatcaccta tggccaaggt agtggtagag cagtttacaa 1020
caacaacatg tacgtcaaca tgtacaacac cgggaatatt gccagagtta acctgaccac 1080
caacacgatt gctgtgactc aaactctccc taatgctgcc tataataacc gcttttcata 1140
tgctaattgt ccttgccaag atattgactt tgcctggtat gagaatggat tgtgggttat 1200
ttattcaact gaagccagca ctggtaacat ggtgattagt aaactcaatg acaccacact 1260
tcagggtgcta aacacttggc ataccaagca gtataaacca tctgcttcta acgccttcac 1320
ggtatgtggg gttctgtatg ccaccctgac tatgaacacc agaacagaag agatttttta 1380
ctattatgac acaaacacag ggaagagggg caaactagac attgtaatgc ataagatgca 1440
ggaaaaagtg cagagcatta actataaccc ttttgaccag aaactttatg tctataacga 1500
tggttacctt ctgaattatg atctttctgt cttgcagaag cccagtaag ctgttttaga 1560
gttaggggtg aagagaaaaa gtttgttgaa aaaatagctc tctccactta cttagatata 1620
tgcaggggtg tctaaaagtg tgttcatttt gcagcaatgt ctagggtgca agttctacca 1680

```



```

cactagagat ctaggacatt tgtcttgatt tggtagagtt tcttgggaat catctgcctc 1740
ttcaggcgca ttttgcaata aagctctgtc agggtagggat tgtcagaggt ctaggggcac 1800
tgtgggccta gtgaagccta ctgtgaggag gcttcactag aagccttaaa ttaggaatta 1860
aggaacttaa aactcagtat ggctcttagg gattctttgt acaggaaaata ttgccaatg 1920
actagtcctc atccatgtag caccactaat tcttccatgc ctggaagaaa cctggggact 1980
tagttaggta gattaatata tggagctcct cgagggaacca aatctccaac ttttttttcc 2040
cctcactaca cctggaatga tgccttgat gtggcagata agtaaatttg gcatgcttat 2100
atattctaca tctgtaaaagt gctgagtttt atggagagag gcctttttat gcattaaatt 2160
gtacatggca aataaatccc agaaggatct gtagatgagg cacttgcttt ttcttttctc 2220
tcattgtcca ccttactaaa agtcagtaga atcttctacc tcataacttc cttccaaagg 2280
cagctcagaa gattagaacc agacttacta accaattcca ccccccacca accoccttct 2340
actgcctact ttaaaaaaat taatagtttt ctatggaaact gatctaagat tagaaaaatt 2400
aatittcttt aatttcatta tggactttta ttacatgac tctaagacta taagaaaatt 2460
tgtggcagt gacaaagtgc tagcatttat tgttatctaa taaagacctt ggagcatatg 2520
tgcaacttat gagtgtatca gttgttgcat gtaatttttg cctttgttta agcctggaac 2580
ttgtaagaaa atgaaaaatt aatttttttt tctaggacga gctatagaaa agctattgag 2640
agtatctagt taatcagtgc agtagttgga aaccttgctg gtgtatgtga tgtgttctg 2700
tgcttttgaa tgactttatc atctagctct tgtctgtttt tctttgatg ttcaagctcc 2760
agtctatagg attggcagtt taaatgcttt actccc 2796

```

<210> 8

<211> 2331

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> (675)

<400> 8

```

tttatcacgg gctcaactgc aacaaaacac ttctttgaca gctccacaaa ctcaggccac 60
agttaggaat caacaatatt ccacagcagc ccagatgcaa gtggaacaac accctcatct 120
gccactcca caacctcagg tcgtggagaa ttacaaacct cagcatcag tccaggctca 180
actgaaataa caacgttacc tggcagtacc acaacaccag gcctcagtga ggcattctacc 240
accttctaca gtagccccag atcaccagac caaacactct cacttgccag catgagaagg 300
tccagcatca gtggagaacc caccagcttg tatagccaag cagagtcaac acacacaaca 360
gcgttccctg ccagcaccac cacctcaggc ctcagtcagg aatcaacaac tttccacagt 420
aagccaggct caactgagac aacactgtcc cctggcagca tcacaacttc atcttttgc 480
caagaattta ccacctcca tagccaacca ggctcagctc tgtcaacagt gtcacctgcc 540
agcaccacag tgccaggcct tagtgaggaa tctaccacct tctacagcag cccaggctca 600
actgaaacca cagcgttttc tcacagcaac acaatgtcca ttcatagtca acaatctaca 660
ccttccctg acagncagg ctteactcac acagtgttac ctgccacct cacaaccaca 720
gacattggtc aggaatcaac agccttccac agcagctcag acgcaactgg aacaacaacc 780
ttacctgcc gctccacagc ctccagacct gttggagaac ctacaacttt ctacatcagc 840
ccatccccta cttacacaac actctttcct gcgagttcca gcacatcagg cctcactgag 900
gaatctacca cttccacac cagtccaagg ttcaacttca caattgtgtc tactgaaagg 960
ctggaaacct tagcaccagg gttgtgccag gaaggacaaa tttggaatgg aaaacaatgc 1020
gtctgtcccc aaggctcagt tggctaccag tgcttgtccc ctctggaatc cttccctgta 1080

```

```

gaaacccccg aaaaactcaa cgcacttta ggtatgacag tgaagtgcac ttacagaaat 1140
ttcacagaaa agatgaatga cgcacccctc caggaatacc agaacttcag taccctcttc 1200
aagaatcgga tggatgtcgt tttgaaggcc gacaatcttc ctacagtatag aggggtgaac 1260
attcggagat tgctcaacgg tagcatcgtg gtcaagaacg atgtcatcct ggaggcagac 1320
tacactttag agtatgagga actgtttgaa aacctggcag agattgtaaa ggccaagatt 1380
atgaatgaaa ctagaacaac tcttcttgat cctgattcct gcagaaaggc catactgtgc 1440
tatagtgaag aggacacttt cgtggattca tcggtgactc cgggctttga cttccaggag 1500
caatgcaccc agaaggctgc cgaaggatat acccagttct actatgtgga tgtcttggt 1560
gggaagctgg cctgtgtgaa caagtgcacc aaaggaacga agtcgcaaat gaactgtaac 1620
ctgggcacat gtcagctgca acgcagtgga ccccgctgc ctgtgoccaa atacgaacac 1680
acactggtac tggggagaga cctgtgaatt caacatcgcc aagagcctcg tgtatgggat 1740
cgtgggggct gtgatggcgg tctgtctgct cgcattgac atccaatca tcttattcag 1800
cctatcccag agaaaaacggc acagggaaca gtatgatgtg cctcaagagt ggcgaagga 1860
aggcaccctt ggcattcttc agaagacggc catctgggaa gaccagaatc tgaggagag 1920
cagattcggc cttgagaacg cctacaacaa ctccggccc acctggaga ctgttgactc 1980
tggcacagag ctccacatcc agaggccgga gatggtagca tccactgtgt gagccaacgg 2040
gggcctccca cctcatctca gctctgttca ggagagctgc aaacacagag cccaccacaa 2100
gcctccgggg cgggtcaaga ggagacgaa gtcaggccct gaagccggtc ctgtcttgag 2160
ctgacagact tggccagtc cctgcctgtg ctctgtctgg ggaaggctgg gggctgtgag 2220
cctctccatc cgggagcttc cagactccca gaagcctcg caccctgtc tctctctggg 2280
tggtcctcca ctctggaatt tccctacca taaaagcaaa tctgaaagct c 2331

```

<210> 9

<211> 909

<212> DNA

<213> Homo sapiens

<400> 9

```

gaggaggtgg ggcaccaacag acaggcgatt aatgcggctc ttaccaggcc aaccaggact 60
acagtataca ttgtggacat tcaggacata gattctgcag ctggggcccg acctcactcc 120
tacctcgatg cctactttgt ctcccccaat gggtcagccc tgaccttga tgagctgagt 180
gtgatgatcc ggaatgatca ggactcgtg acgcagctgc tgcagctggg gctgggtggtg 240
ctgggctccc aggagagcca ggagtcagac ctgtcgaaac agctcatcag tgtcatcata 300
ggattgggag tggctttgct gctggtcctt gtgatcatga ccatggcctt cgtgtgtgtg 360
cggaagagct acaaccggaa gcttcaagct atgaaggctg ccaaggaggc caggaagaca 420
gcagcagggg tgatgccctc agcccctgcc atcccaggga ctaacatgta caaactjag 480
cgagccaacc ccatgtctga cctccccaac aaagacctgg gcttggagta cctctctccc 540
tccaatgacc tggactctgt cagcgtcaac tccctggacg acaactctgt ggatgtggac 600
aagaacagtc aggaaatcaa ggagcacagg ccaccacaca caccaccaga gccagatcca 660
gagccctga gcgtggtcct gttaggacgg caggcaggcg caagtggaca gctggagggg 720
ccatcctaca ccaacgctgg cctggacacc acggacctgt gacagggggc cccactcttc 780
tggacctctt gaagaggccc taccacccc taactgcacc tgtctcctg gagatgaaaa 840
tatatgacgc tgccctgcct cctgcttttg gccaatcacg gcagacaggg gttggggaaa 900
tattttatt
909

```

<210> 10

<211> 510

<212> PRT

<213> Homo sapiens

<400> 10

Met Arg Pro Gly Leu Ser Phe Leu Leu Ala Leu Leu Phe Phe Leu Gly
 1 5 10 15

Gln Ala Ala Gly Asp Leu Gly Asp Val Gly Pro Pro Ile Pro Ser Pro
 20 25 30

Gly Phe Ser Pro Phe Pro Gly Val Asp Ser Ser Ser Ser Phe Ser Ser
 35 40 45

Ser Ser Arg Ser Gly Ser Ser Ser Arg Ser Leu Gly Ser Gly Gly
 50 55 60

Ser Val Ser Gln Leu Phe Ser Asn Phe Thr Gly Ser Val Asp Asp Arg
 65 70 75 80

Gly Thr Cys Gln Cys Ser Val Ser Leu Pro Asp Thr Thr Phe Pro Val
 85 90 95

Asp Arg Val Glu Arg Leu Glu Phe Thr Ala His Val Leu Ser Gln Lys
 100 105 110

Phe Glu Lys Glu Leu Ser Lys Val Arg Glu Tyr Val Gln Leu Ile Ser
 115 120 125

Val Tyr Glu Lys Lys Leu Leu Asn Leu Thr Val Arg Ile Asp Ile Met
 130 135 140

Glu Lys Asp Thr Ile Ser Tyr Thr Glu Leu Asp Phe Glu Leu Ile Lys
 145 150 155 160

Val Glu Val Lys Glu Met Glu Lys Leu Val Ile Gln Leu Lys Glu Ser
 165 170 175

Phe Gly Gly Ser Ser Glu Ile Val Asp Gln Leu Glu Val Glu Ile Arg
 180 185 190

Asn Met Thr Leu Leu Val Glu Lys Leu Glu Thr Leu Asp Lys Asn Asn
 195 200 205

Val Leu Ala Ile Arg Arg Glu Ile Val Ala Leu Lys Thr Lys Leu Lys
 210 215 220

Glu Cys Glu Ala Ser Lys Asp Gln Asn Thr Pro Val Val His Pro Pro
 225 230 235 240

Pro Thr Pro Gly Ser Cys Gly His Gly Gly Val Val Asn Ile Ser Lys
 245 250 255

Pro Ser Val Val Gln Leu Asn Trp Arg Gly Phe Ser Tyr Leu Tyr Gly
 260 265 270

Ala Trp Gly Arg Asp Tyr Ser Pro Gln His Pro Asn Lys Gly Leu Tyr
 275 280 285

Trp Val Ala Pro Leu Asn Thr Asp Gly Arg Leu Leu Glu Tyr Tyr Arg
 290 295 300

Leu Tyr Asn Thr Leu Asp Asp Leu Leu Leu Tyr Ile Asn Ala Arg Glu
 305 310 315 320

Leu Arg Ile Thr Tyr Gly Gln Gly Ser Gly Thr Ala Val Tyr Asn Asn
 325 330 335

Asn Met Tyr Val Asn Met Tyr Asn Thr Gly Asn Ile Ala Arg Val Asn
 340 345 350

Leu Thr Thr Asn Thr Ile Ala Val Thr Gln Thr Leu Pro Asn Ala Ala
 355 360 365

Tyr Asn Asn Arg Phe Ser Tyr Ala Asn Val Ala Trp Gln Asp Ile Asp
 370 375 380

Phe Ala Val Asp Glu Asn Gly Leu Trp Val Ile Tyr Ser Thr Glu Ala
 385 390 395 400

Ser Thr Gly Asn Met Val Ile Ser Lys Leu Asn Asp Thr Thr Leu Gln
 405 410 415

Val Leu Asn Thr Trp Tyr Thr Lys Gln Tyr Lys Pro Ser Ala Ser Asn
 420 425 430

Ala Phe Met Val Cys Gly Val Leu Tyr Ala Thr Arg Thr Met Asn Thr
 435 440 445

Arg Thr Glu Glu Ile Phe Tyr Tyr Tyr Asp Thr Asn Thr Gly Lys Glu
 450 455 460

Gly Lys Leu Asp Ile Val Met His Lys Met Gln Glu Lys Val Gln Ser
 465 470 475 480

Ile Asn Tyr Asn Pro Phe Asp Gln Lys Leu Tyr Val Tyr Asn Asp Gly
 485 490 495

WO 99/60161

PCT/US99/10498

Tyr Leu Leu Asn Tyr Asp Leu Ser Val Leu Gln Lys Pro Gln
500 505 510

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/10498

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C12Q 1/68; G01N 33/53, 33/574

US CL : 435/4, 6, 7.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/4, 6, 7.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Scisearch, HCAPLUS, medline, biosis, membase, wpids, jicst-eplus, biobusiness, biotechds, phin, phic, embal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96/39419 A1 (HUMAN GENOME SCIENCES, INC.) 12 December 1996, see entire document.	1-5

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

16 AUGUST 1999

Date of mailing of the international search report

10 SEP 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

NANCY JOHNSON

Telephone No. (703) 308-0196

Form PCT/ISA/210 (second sheet)(July 1992)*

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/10498

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☒ Claims Nos.: 6
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

The sequence listing in computer readable form did not comply with required standards. The disc could not be preprocessed. Thus, a search of claim 6, drawn to SEQ ID NO:3, 4, 5 or 7, could not be carried out.

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet(1))(July 1992)*